



Star of the Sea Primary School

Dealing with Head Lice

School Based Policy & Procedure

Originally written: 2014

Date for Review: 2017

Head Lice is a normal and quite a frequent occurrence in all schools and no doubt causes some consternation when it happens and especially when it is a returning issue.

Star of the Sea Primary School follows the following procedure to support families in dealing with a case of head lice whilst respecting the dignity and privacy of those concerned.

For the concern of all, our School Uniform Policy states "For health reasons, hair must be off the collar. Children with hair that is collar length or longer are required to have hair fully tied up, back off the face. Plaits, braids and buns are recommended to minimize transference of lice."

Other prevention tips are also included in the information provided by the Department of Health at the conclusion of this policy.

Procedures

Children will be excluded from school until treatment has been commenced. Where possible exclusion occurs at the end of the school day, unless otherwise deemed necessary.

When a case of head lice is noticed in class, families are notified so that they may check their child's hair to minimize the spread.

The child who has been identified as having head lice is sent a separate notice advising a treatment is necessary and information regarding head lice and treatment methods are also made available from the Health Department. A child may return to school when all live head lice have been removed. A few remaining eggs is not a reason for continued exclusion.

Children should continue to have their hair checked daily for up to 10 days. All eggs will need to be removed. Other household members should also be checked.

In order to help each other we will introduce an occasional "Nits Blitz" and when that is called all families will be asked to check their children's hair to further minimise the spread.

The School Health Nurse will be asked to provide further assistance to the school and families in the case of repeated incidences.



Star of the Sea

CATHOLIC PRIMARY SCHOOL

(Insert Date)

Dear Parents

An incidence of HEAD LICE has been detected in (insert class).

We ask all parents to please inspect their child/all family members' hair **tonight** then daily for one week and commence treatment if necessary.

Should head lice be detected, children are to be excluded from school until appropriate treatment has been undertaken.

Further information about head lice can be found at www.healthywa.wa.gov.au/Healthy-WA/Articles/F_I/Head-lice

Thank you for your cooperation in this matter.

Yours faithfully,

Assistant Principal



Star of the Sea

CATHOLIC PRIMARY SCHOOL

NOTIFICATION OF HEAD LICE

Dear Parents

(student name) was found to have head lice today.

You are required to treat (his/her) tonight and then check daily for up to 2 weeks. Other household members should also be checked.

An information pamphlet has been sent home along with this note. Your child may return to school following effective treatment. Please sign the slip below and present to the school teacher on your child's return to school.

Please note: If your child has received recent treatments please note date of treatment below and remember if some eggs remain in the hair they will need to be removed. Also follow up treatment may be necessary.

Thanking you in anticipation of your co-operation in keeping this problem under control.

Yours faithfully

Assistant Principal

✂ _____

My child _____ has been treated for head lice with a pharmacy approved lotion.

Signed _____ Date _____

Head lice

What are head lice?

Head lice are tiny insect parasites that live on your head and feed on your scalp (the skin covering your head).

They reproduce by laying their eggs (nits) on your hair shaft (the part of your hair closest to the scalp).

Head lice are not dangerous, do not carry diseases, and are not a sign of poor hygiene (cleanliness).

How do you get head lice?

Head lice are spread by head-to-head contact with another person who has head lice. This kind of contact includes doing group work at school, playing, or hugging.

Head lice can run from one head to another in seconds. Head lice cannot fly, jump or swim, but they can sometimes swing from one hair to another.

Brushes and combs are unlikely to transfer head lice or their eggs, as these are very hard to detach from the hair shaft.

Head lice are not spread through bed linen, clothing or head gear (hats and helmets) as they do not leave the scalp unless they are moving to another scalp, or unless they are dead or dying.

Signs and symptoms

Your scalp may itch as your skin reacts to the saliva of the head lice. This itchiness can take weeks to develop. If you have had head lice before, your skin may become less sensitive and there may be little or no itch.

You may also be able to see head lice crawling in your hair, although they can be difficult to spot as they move quickly.

- Adult lice are usually dark brown and about 2 to 3 mm long.
- Hatchlings (young lice) are often a lighter brown colour and about 1 to 2 mm long.
- Eggs will be attached to the hair shaft. They can be very tiny and hard to see, especially newly-laid eggs close to the scalp. They are grey-white and about the size of a grain of salt.

Checking for head lice

Unless you can easily see the head lice, the speed at which they can move makes checking dry hair unreliable.

To check hair for head lice:

- Carefully comb plenty of hair conditioner through dry hair. The conditioner slows the head lice down so they can be trapped in the comb.
- Comb hair again in sections using a metal fine-tooth 'nit comb' (available from most chemists).
- Wipe the waste from the comb on a white paper towel. Look for head lice and eggs using a magnifying glass in strong light, such as sunlight.
- Use a magnifying glass in strong light to examine hair close to the scalp, especially behind the ears and at the back of the neck.

Distract children while you check for head lice by offering them a favourite video or game, a book, or a head massage.

What can be mistaken for head lice?

Other objects in the hair can be mistaken for head lice or eggs. These include sand, dandruff, flakes of hair spray, ants, aphids, or other small insects.

If you find head lice

- Check to see if anyone else living in your house has head lice – use the hair conditioner method described above.
- Choose the treatment method that best suits you and your family and follow the instructions carefully.
- Inform your school, child care centre, playgroup, neighbours and relatives so that other children and adults who have been in close contact can be checked.
- When the treatment is completed and all head lice and eggs have been removed, continue to check hair weekly for further outbreaks.

Stay home from school

Under the *School Education Act 1999*, if your child has head lice the principal may keep him or her away from school until treatment has started.

Your child may return to school when all live head lice have been removed. There is no need to stay away from school if there are only a few remaining eggs, but you must continue treatment over the following 10 days to ensure that all eggs and hatchlings have been removed.

Prevent further outbreaks

- Check hair regularly for head lice and eggs. Early treatment of head lice can prevent it spreading further to close contacts.
- Tie long hair back.
- Hair gel or mousse may help keep stray hair strands from coming in contact with other heads.
- Avoid sharing combs or brushes.
- If you have recently had head lice, leave brushes and combs in hot water for 10 minutes after use to kill any trapped head lice or eggs. Clean them with an old toothbrush to help remove any waste.
- Remind your child to avoid head-to-head contact with other children, for example when working at the computer with others, playing or hugging.

This publication is provided for education and information purposes only. It is not a substitute for professional medical care. Information about a therapy, service, product or treatment does not imply endorsement and is not intended to replace advice from your healthcare professional. Readers should note that over time currency and completeness of the information may change. All users should seek advice from a qualified healthcare professional for a diagnosis and answers to their medical questions.